

# **Geological and Geophysical Support for the “Geoclutter” Acoustic Reconnaissance Field Program (Phase I)**

John A. Goff  
James A Austin, Jr.  
Institute for Geophysics, University of Texas at Austin  
4412 Spicewood Springs Rd., Bldg. 600, Austin, TX 78759  
phone: (512) 471-0476 fax: (512) 471-0999 email: [goff@utig.ig.utexas.edu](mailto:goff@utig.ig.utexas.edu)

Award Number: N00014-99-1-1046  
<http://www.ig.utexas.edu/staff/goff/goff.html>

## **LONG-TERM GOALS**

Identify and characterize, sufficient for acoustic modeling purposes, seafloor and subseafloor structures that may give rise to signal-like returns (“geoclutter”) on long-range active sonar systems.

## **OBJECTIVES**

The primary objective of this work is to provide geologic and geophysical support for the planned Geoclutter acoustic reconnaissance experiment (Phase I. A. of the overall program; see below), to be conducted on the New Jersey shelf south of the Hudson Shelf Valley (Figure 1). This will involve: (1) preparing an environmental assessment; (2) identifying likely geoclutter targets based on existing bathymetry, seismic reflection, and sampling data; (3) making available digital maps and seismic interpretation profiles for use in acoustic modeling and cruise preparation; (4) participation in the acoustic reconnaissance cruise; and (5) initial preparations for a forthcoming geophysical survey of acoustic “hotspots” that are identified in the reconnaissance.

## **APPROACH**

The premise for this work is that, in any littoral area, buried geologic features can contribute significantly to acoustic reverberation, which affects tactical ASW sonar systems. Proper acoustic processing, coupled with quantitative geologic models, can be used to distinguish these buried features from real (man-made) targets. Complexity arises from STRATAFORM studies on the continental shelf off New Jersey that have shown the general lack of predictability of the shallow subsurface using seafloor imagery, even with 100% coverage.

The primary goal of the Geoclutter program will be to assess geologic clutter/reverberation issues in a well-characterized shallow-water environment. The mid-outer continental shelf off New Jersey provides an opportunity, because both bathymetry (a known and prominent cause of backscatter) and the shallow subsurface have been mapped in detail as a result of STRATAFORM. The Geoclutter program will consist of three field program phases: (I) a monostatic acoustic reconnaissance of the New Jersey shelf region to identify acoustic “hot spots” for further study; (II) detailed geologic and geophysical survey of the hot spots identified in Phase I; and (III) a full bistatic acoustic experiment

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>30 SEP 1999</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-1999 to 00-00-1999</b>	
4. TITLE AND SUBTITLE <b>Geological and Geophysical Support for the 'Geoclutter' Acoustic Reconnaissance Field Program (Phase I)</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>University of of Texas at Austin, Institute for Geophysics (UTIG, 4412 Spicewood Springs Road, Bldg. 600, Austin, TX, 78759</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>5</b>	19a. NAME OF RESPONSIBLE PERSON
a REPORT <b>unclassified</b>	b ABSTRACT <b>unclassified</b>	c THIS PAGE <b>unclassified</b>			

focusing on the chosen areas. The focus of our present work is primarily on the Phase I experiment. Dr. N. Makris of MIT is the PI in charge of the Phase I experiment and preliminary acoustic modeling.

Our initial efforts have been aimed at identifying potential geoclutter target sites for the purpose of planning the Phase I experiment, and providing to Dr. Makris profiles of seafloor and shallow stratigraphic morphology. There are a number of candidate clutter targets that can be identified in bathymetry, sidescan and seismic reflection data that presently exist in the proposed survey area. These include: buried channels with highly reflective channel walls and thalwegs from 2D and 3D Huntet profiling; iceberg scour marks, erosion pits, large sand ridges, surface or near-surface outcroppings of seismically reflective subsurface horizons, and areas of highly reflective, probably gravelly, seafloor identified from the sidescan.

## **WORK COMPLETED**

We have completed a preliminary map of potential geoclutter target sites within the STRATAFORM New Jersey natural laboratory (Figure 2) and provided to Dr. Makris a profile of bathymetry and subsurface stratigraphy for use in his preliminary modeling work (Figure 3).

## **RESULTS**

The results of this work thus far are principally contained in Makris' progress report. One of the more salient impacts of the acoustic modeling work based on our profiles has been to recognize the critical importance of accurate measurement of seafloor sediment properties, especially sound velocity. Given the extremely low grazing angles associated with the operation of long range active sonar systems in shallow water, small changes in the sound velocity of seafloor sediments can have a dramatic impact on the level of acoustic energy that is transmitted into the subsurface.

## **IMPACT/APPLICATIONS**

The newly formed ONR-sponsored "Geoclutter" program is a ~5-year field and analysis effort: (1) to understand, characterize, and predict lateral and vertical, naturally-occurring heterogeneities that may produce discrete acoustic returns at low grazing angles in a mid-outer shelf test site off the U.S. (New Jersey), and then (2) to conduct precise acoustic reverberation experiments at this site to understand, characterize, and potentially mitigate the geologic clutter, so that the false alarms, or detects, of tactical sonar systems encountered in this marine geologic environment around the world can be characterized properly. The goal of the planned field experiments off the U.S. east coast is to understand the process of acoustic reverberation from the seabed in shallow water, with the objective of designing physics-based signal processing ("detection") algorithms to distinguish these naturally-occurring features on the world's continental shelves (e.g., shallowly-buried meandering channels, surface outcrop of reflective horizons, erosion scars) from man-made targets of similar dimensions (e.g., submarines).


The products that we are generating under this grant will greatly facilitate the planning and operation of the Phase I Geoclutter acoustic reconnaissance field program. This work also constitutes preliminary step for the Phase II work of detailed geological and geophysical survey of the target sites that will ultimately be chosen for intensive acoustic study in Phase III.

## TRANSITIONS


The products that we are generating are being used by Dr. Makris for the planning of the Phase I acoustic reconnaissance experiment and for preliminary acoustic modeling work.

## RELATED PROJECTS


The New Jersey shelf was chosen as the focus for the Geoclutter work in large part because of the intensive data collection that has already occurred there as part of both the ONR STRATAFORM program and ODP site survey work.



**Figure 1. Location figure for Geoclutter experiment. Contour depth in meters. Offshore shaded region is area covered by STRATAFORM New Jersey swath sonar data.**



**Figure 2. Possible geoclutter targets identified from swath map and ultra-high resolution seismic reflection data. See Figure 1 for location. The STRATAFORM swath bathymetry data are displayed with artificial illumination from the north. Contours in meters are from regional data compiled by NGDC. Light blue lines indicate tracklines for available Hunttec ultra-high resolution seismic reflection data. Three different types of possible geoclutter targets are identified: red ovals indicate bathymetric targets; yellow lines indicate important changes in sedimentary properties at or near the seafloor, as inferred from backscatter data or by outcrops of prominent subsurface reflectors; black dashed lines indicate shallowly buried (< 10 m) channels (see Figure 3).**



***Figure 3. Seismic reflection interpretation profile on the New Jersey shelf indicating both seafloor and subsurface complexities that could give rise to "geoclutter" signals. This profile has been provided to Makris for his preliminary modeling work.***